

The Impact of the NACRE Compilation on the Big Bang Nucleosynthesis

Elisabeth Vangioni-Flam^a, Alain Coc^b and Michel Cassé^{c a}

^aInstitut d'Astrophysique de Paris, 98^{bis} Bd Arago 75014 Paris, France

^b CSNSM, IN2P3-CNRS-UPS, Bâtiment 104, 91405 Orsay Campus, France

^c SAp, DAPNIA, DSM, CEA, Orme des Merisiers, 91191 Gif sur Yvette CEDEX France

We update the Big Bang Nucleosynthesis (BBN) calculations on the basis of the recent NACRE compilation of reaction rates. We estimate the uncertainties related to the nuclear reaction rates on the abundances of D , ${}^3\text{He}$, ${}^4\text{He}$, ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^9\text{Be}$, ${}^{10}\text{B}$ and ${}^{11}\text{B}$ of cosmological and astrophysical interest. We use lithium as the main indicator of the baryon density of the Universe, rather than deuterium.

1. Introduction

Recently, the new NACRE compilation of thermonuclear rates[1] has been published and observations of light isotopes have flourished. It is thus timely to reassess the determination of the baryonic density of the Universe in the light of advances in nuclear physics and astronomical observations. To calculate primordial abundances up to B , the BBN network is constituted by about 60 reactions[2] out of which, 22 are included in the NACRE compilation. The main improvements with respect to former compilations concern a better traceability to the source of nuclear data and the availability of upper and lower rate limits. The primordial abundances of the light elements D , ${}^{3,4}\text{He}$ and ${}^7\text{Li}$ are governed by the expansion rate of the Universe and the cooling it induces. Under the classical assumptions, these abundances depend only on the baryon to photon ratio η , related to the baryonic parameter by $\eta_{10} = 273.\Omega_B.h^2$ with $h = H/100$ km/s.Mpc.

2. Discussion

D/H is measured in three astrophysical and/or cosmological sites : local interstellar medium : 0.5 to $2. \cdot 10^{-5}$ [3,4], protosolar nebula : $3.\pm 0.3 \cdot 10^{-5}$ [5], absorbing clouds on the sightlines of quasars : $3.$ to $4. \cdot 10^{-5}$, [6] and 0.8 to $3. \cdot 10^{-4}$ [7]. The primordial abundance of ${}^4\text{He}$, Y_p is measured in low metallicity extragalactic HII regions and in HII regions in the Small Magellanic Cloud. Two ranges emerge: $Y_p = 0.238 \pm 0.004$ [8,9] and $Y_p = 0.245 \pm 0.004$ [10]. Determination of Li in halo stars[11–13] indicate that the Spite plateau is exceptionally thin (< 0.05 dex). We adopt the following range: $1.5 \cdot 10^{-10} < {}^7\text{Li}/H < 2.2 \cdot 10^{-10}$, taking into account a maximum destruction of 0.1 dex. The ${}^6\text{Li}$ abundance has also been determined in two halo stars[14–16]. We confirm that BBN calculated abundances of ${}^9\text{Be}$, ${}^{10}\text{B}$ and ${}^{11}\text{B}$ are negligible with respect the measured ones in the more metal




Figure 1. Theoretical primordial abundances of ${}^4\text{He}$ (by mass), D , ${}^3\text{He}$ and ${}^7\text{Li}$ (by number) vs the baryon/photon ratio, η , using the reaction rates from the NACRE compilation[1]. Solid lines: using recommended values of the reaction rates. Dashed lines: using high or low values of the reaction rates[2] with possible error compensation. Dash-dotted lines: most extreme results following all combination of high and low rates. Horizontal lines indicate the error bands related to different observations.

poor stars. The uncertainty on the $\text{D}(\alpha, \gamma){}^6\text{Li}$ rate, and hence on of ${}^6\text{Li}$ production, is high. At the upper limit, it could lead to a primordial ${}^6\text{Li}$ abundance within reach of future observations, to be used as a new cosmological constraint. Anyway, the bulk of the abundances of these elements can be explained in term of spallation of fast carbon and oxygen in the early Galaxy[17]. Considering ${}^7\text{Li}$ observational constraints (see Fig. 1), in the most favorable case where errors compensate[2] two possible ranges emerge: *i*) $1.5 < \eta_{10} < 1.9$ corresponding to $0.013 < \Omega_B < 0.019$ ($h_{100} = 0.65$) in good concordance with the error boxes related to the observed high D and low ${}^4\text{He}$ *ii*) $3.3 < \eta_{10} < 5.1$ corresponding to $0.029 < \Omega_B < 0.045$ in fair agreement with a low D/H and a high ${}^4\text{He}$. In the most pessimistic case, where errors are maximized (considering all combinations of high and low NACRE rates, see Fig. 1) the η_{10} ranges are wider (1.2 to 2) and (3 to 5.1). Comparing the baryonic density to that of luminous matter ($0.002 < \Omega_L < 0.004$ [18]) in the Universe shows the necessity of baryonic dark matter. The observations of the Lyman alpha forest clouds between the redshifts 0 and 5, lead to a corresponding

$\Omega_B = 0.03 \pm 0.01$, taking into account the uncertainty related to ionized hydrogen[19]. This value is thought to reflect the bulk of the baryons at large scale.

3. Conclusion

Big bang nucleosynthesis deserves permanent care since it gives access to the baryon density which is a key cosmological parameter. This work has been aimed at integrating the last development in both fields of nuclear physics and observational abundance determination of light elements. The update of the reaction rates of the BBN using the NACRE compilation has been made. ${}^6\text{Li}$ is affected by the large uncertainty of the $D(\alpha, \gamma){}^6\text{Li}$ reaction. A refined measurement of this reaction is of great cosmological interest. Owing to the high observational reliability of the halo star ${}^7\text{Li}$ abundance data with respect to the D data available which are both rare and debated, we choose it as the leading baryometer. Due to the valley shape of the Li curve, we get two possible η ranges which are consistent respectively with a high D and low ${}^4\text{He}$ observed values and with a low D and high ${}^4\text{He}$ observed values. At present, none of these solutions can be excluded.

REFERENCES

1. C. Angulo, M. Arnould, M. Rayet et al., Nucl. Phys. A656 (1999) 3.
2. E. Vangioni-Flam, A. Coc & M. Cassé, Astron. Astrophys. 360 (2000) 15.
3. J. Linsky et al., Astrophys. J. 451 (1995) 335.
4. A. Vidal-Madjar, in 'The light Elements and their Evolution', IAU Symp. 198, ASP Conf. Series., Edts L. da Silva, R. de Meideros and M. Spite, (2000) in press
5. A. Drouart, B. Dubrulle, D. Gautier & F. Robert, Icarus 140 (1999) 129.
6. D. Tytler et al., in 'Nobel Symposium 109: Particle Physics and the Universe', Physica Scripta T85 (2000) 12.
7. J.K. Webb et al., Nature 383 (1997) 250.
8. B. Fields & K. Olive, Astrophys. J. 506 (1998) 177.
9. M. Peimbert & A. Peimbert, in 'The light Elements and their Evolution', IAU Symp. 198, ASP Conf. Series. Edts L. da Silva, R. de Meideros and M. Spite, (2000) in press.
10. Y.I. Izotov & T.X. Thuan, Astrophys. J. 500 (1998) 1888.
11. M. Spite, P. Francois, P.E. Nissen & F. Spite, Astron. Astrophys. 307 (1996) 172.
12. P. Bonifacio & P. Molaro, Mon. Not. R. Astron. Soc. 285 (1997) 847.
13. S.G. Ryan, J.E. Norris & T.C. Beers, Astrophys. J. 523 (1999) 654.
14. L.M. Hobbs & J.A. Thorburn, Astrophys. J. 491 (1997) 772.
15. V.V. Smith, D.L. Lambert & P.E. Nissen, Astrophys. J. 506 (1998) 405.
16. R. Cayrel et al., Astron. Astrophys. 343 (1999) 923.
17. E. Vangioni-Flam, M. Cassé & J. Audouze, Physics Report 333-334 (2000) 365.
18. P. Salucci & M. Persic, Mon. Not. R. Astron. Soc. 309 (1999) 923.
19. R. Riedeger, P. Petitjean & J.P. Mucket, Astron. Astrophys. 329 (1998) 30.